

Slobber-Mash with extra Linol

PRODUCT DESCRIPTION

Slobber-Mash is a very palatable, coarse mixture of flaked cereals, corn, toasted linseed and wheat bran, completed with extra vitamins and minerals.

Additionally, it contains extra linoleic acids, linseed oil and sunflower seeds.

Slobber-Mash provides support during the moult in autumn and spring. Also recommended for hard-keepers or horses that have difficulties in maintaining or increasing bodyweight.

- Easily absorbable and easily digestible
- For extra shiny coat
- Supports a healthy bowel movement
- Ideal for older horses, hard-keepers and horses in poor condition
- Can be fed both dry or soaked with water



Nutrient /kg					
VEP	925	Vit. A	18.750 IE	Iron	326mg
Carbohydrates	53%	Vit. D-3	3.000 IE	Copper	24mg
Starch	38,2%	Vit. E	330mg	Zinc	264mg
Sugar	5,6%	Biotin	167mcg	Manganese	123mg
Crude fibre	7,6%	Vit. K3	1,4mg	Cobalt	2,0mg
Dig. protein	7,9%	Vit. B1	4,1mg	Iodine	1,5mg
Crude protein	10,5%	Vit. B2	6,9mg	Selenium	426mcg
Crude fat	4,0%	Vit. B6	3,7mg		
Crude ash	7,1%	Vit. B12	27,5mcg	Lysin	3,8g
Calcium	11,9gr	Panthenic acid	19,4mg	Meth+cyst	3,9g
Phosphorous	4,2gr	Niacin	38mg	Threonin	3,6g
Magnesium	2,4gr	Folic acid	4,1mg	Tryptofan	1,2g
Sodium	2,5gr	Cholin-chloride	206mg		
Potassium	9,4gr				
Packaging: 20 kg /bag					

Datasheet subject to change April 2019

Due to its specific composition the Slobber-Mash works very stimulating to the skin, hair and Hoofs.

Result: a shiny appearance!

FEEDING GUIDELINE

Best to be used with water (but can also be fed dry). Soak 1 part of Slobber- Mash with 1 part of water for 15-20 minutes. This way, it can be absorbed easily, also by (older) horses with missing teeth.

It is important to feed a diet and quantity that meets the lifestyle and needs of the horse. A general guideline for the recommended total amount of concentrates per day:

- For maintenance and basic sport: 0.2 to 0.5 kg per 100 kg bodyweight.
- For (top) sport: 0.5 to 1.0 kg per 100 kg bodyweight.

Divide this daily ration into at least 2-3 meals per day. Always try to feed a minimum of 1.5% of bodyweight as good quality forage. Fresh drinking water must be available at all times.